

Package: SeedImbibition (via r-universe)

September 18, 2026

Type Package

Title Seed Imbibition Percentage

Version 0.1.0

Maintainer Tanuj Misra <tanujmisra102@gmail.com>

Description Imbibition causes seeds to expand, which results in the seed coat or testa being broken. Seed germination begins with imbibition. Imbibition aids in the transport of water into the developing ovules. Imbibition is required during the first stages of root water absorption.

License GPL-3

Encoding UTF-8

NeedsCompilation no

Author Sudip Kumar Dutta [aut, cph], Tanuj Misra [aut, cre, cph]

Repository <https://tanujrbcu.r-universe.dev>

Date/Publication 2024-06-01 02:44:57 UTC

RemoteUrl <https://github.com/cran/SeedImbibition>

RemoteRef HEAD

RemoteSha afded3e652fd35b45bf215ec9ec36c4c24a451c4

Contents

SeedImbibition	2
Index	3

SeedImbibition	<i>Seed Imbibition Percentage</i>
----------------	-----------------------------------

Description

Imbibition causes seeds to expand, which results in the seed coat or testa being broken. Seed germination begins with imbibition. Imbibition aids in the transport of water into the developing ovules. Imbibition is required during the first stages of root water absorption. Imbibition percentage of seeds was calculated using the following formula (Govender et al., 2008) <doi:10.1016/j.sajb.2007.10.006>

Usage

```
seed_imbib(wt_of_seed_after_imbib, initial_wt_of_seed)
```

Arguments

```
wt_of_seed_after_imbib
                        Weight of seeds after imbibition
initial_wt_of_seed
                        Initial weight of seeds
```

Value

Seed Imbibition Percentage

References

Govender, V., Aveling, T.A.S., Kritzing, Q., 2008. The effect of traditional storage methods on germination and vigour of maize (*Zea mays* L.) from northern KwaZulu-Natal and southern Mozambique. *S. Afr. J. Bot.* 74, 190–196. \ Dutta, S. K., Layek, J., Akoijam, R. S., Boopathi, T., Vanlalhmangaiha, Saha, S., & Prakash, N. (2019). Seaweed extract as natural priming agent for augmenting seed quality traits and yield in *Capsicum frutescens* L. *Journal of applied phycology*, 31, 3803-3813.

Examples

```
wt_of_seed_after_imbib <- 2.5
initial_wt_of_seed <- 1.7
seed_imbib(wt_of_seed_after_imbib, initial_wt_of_seed)
```

Index

* **Seed Imbibition Percentage**

SeedImbibition, [2](#)

seed_imbib (SeedImbibition), [2](#)

SeedImbibition, [2](#)